

Shoulder Arthroplasty | Physical Therapy Protocol

At UAMS | Health and at **Train • Recover • Move**, our goal is to create an environment that is safe for the healing structures, exciting for the patient, and able to provide an open and transparent line of communication with the therapist. If you ever have any questions or concerns, please feel free to give us a call and we would be more than happy to discuss any concerns or questions you might have.

Our protocol can be divided into three distinct and separate phases:

Phase One (Day 1-Week 4-6): Protect

Phase Two (Week 6-Week 12): Mobility

Phase Three (Week 12+): Strength

Before we dive into the details of the separate phases, we would love to take some time answer some common questions our patients often ask considering this surgery and rehabilitation process:

Do I really need to wear my sling?

- Based on the size and severity of your rotator cuff tear, you will be instructed to wear your sling for the first four to six weeks following your surgery. Unwanted movement early on can hinder tissue that can lead to an unstable shoulder.
- We encourage the patient to wear the sling at all times. There are components on the sling that might seem bulky, but we need you to keep these foam cushions in the correct places for the entire four to six weeks unless notified by your surgeon.
- You may remove the sling only for the exercises that has been prescribed by your physical therapist, for using ice, and for dressing or showering.

When do I begin Physical Therapy?

- When physical therapy begins will depend on the type of arthroplasty that you receive and the size of the components.
- **The frequency and when a patient starts physical therapy can be adjusted by the surgeon.**

How often do I need to go to physical therapy?

- The frequency of physical therapy will typically start at two-three times a week. This frequency can also be adjusted by your surgeon or your physical therapist's recommendations.
- Once strength training begins, the patient's frequency can be reduced to one or two times a week considering good home exercise compliance.

When can I start strengthening?

- We do not recommend strength training until approximately week twelve after surgery.

Well, when do I get back to sports and my recreational activities?

- Sports that demand high level use of the upper extremity (examples: baseball, volleyball, football, and tennis) will typically be asked to wait six months before returning to the sport or the recreational activity.
- Please seek permission from your surgeon and physical therapist before starting back to any sporting or recreational activity.

Well, when do I get back to sports and my recreational activities?

- Sports that demand high level use of the upper extremity (examples: baseball, volleyball, football, and tennis) will typically be asked to wait six months before returning to the sport or the recreational activity.
- Please seek permission from your surgeon and physical therapist before starting back to any sporting or recreational activity.

Is there anything I need to avoid after surgery?

- We encourage any patient who smokes to avoid smoking after surgery for the first twelve weeks. Smoking can correlate with delayed and improper tissue healing.
- Avoid movements such as reaching up your back or moving your arm away from your body during the first four weeks. If you are at a computer, keep your elbow by your side at all times during the initial six weeks.
- No pushing, pulling, or lifting anything larger than a cup of coffee (approx. 1-2 pounds). Avoid pushing yourself up from a chair, bed, or from your car for the initial six weeks following your surgery.
- We do not want our patient to take any kind of anti-inflammatories (naproxen sodium, ibuprofen, or Aleve) for the first twelve weeks following surgery. In order to manage any pain, we encourage our patients to use their ice machine as needed.

I have friends who have had a shoulder replacement, will my experience be like theirs?

- There are several factors that are specific to the patient that will affect their recovery. Some of these factors can include the type of replacement, the surgical approach, the quality of the tissue, the quality of the remaining tissue, and the mechanism of failure. Each patient's recovery is very individualized. It will generally take approximately four six months before we hear patients say they feel "back to normal." However, for other patients it might take up to 1 year before full recovery takes place. any pain, we encourage our patients to use their ice machine as needed.

PHASE ONE

Day 1 to Week 4-6 | Protect

GOALS:

Protect surgical repair by avoiding unwanted strain to insertion site

Maintain elbow, hand, and wrist mobility

Improve neuromuscular activation that's been inhibited by edema and pain

Normalize scapular position and posture

Address cervical and thoracic posture / mobility

THINGS TO AVOID:

There should be no active and aggressive internal rotation movement of the shoulder.

No pushing, pulling, or lifting anything larger than a cup of coffee (approx. 1-2 pounds).

Avoid pushing yourself up from a chair, bed, or from your car.

While we encourage you to tease, touch, and nudge any pain you might experience, please don't push through the pain. Do not be aggressive with any passive mobility that might cause muscle spasms or increased sharp pains in the shoulder.

WEEK THREE:

Begin physical therapy 1x / week.

Address portal site mobility, drainage, excessive redness, or discoloration

Keep any motion in scapular plane in order to stay in open packed position

Patient positioning in supine or reclined - avoid supine or prone positions

If aquatic option is available, this can be utilized if incision site is completely healed

- Scapular plane passive movement only to 90°
- Avoid internal rotation
- No external rotation greater than 30°

Begin home exercise program that is to be completed 2-3 times each day:

- Pendulums
- Elbow, hand, and wrist active mobility exercises. No weights
- Submaximal and PAIN FREE isometrics
 - * Flexion, extension, abduction, adduction, and external rotation
- Scapular squeezes, depression, and rolls
- Passive scapular plane elevation to 90° with pulley apparatus
- Ice and modalities as needed
 - * No motoric response with electrical stimulation

WEEK FOUR:

Continue to monitor incision healing progression

Assess patient compliance / understanding with home exercise program

Patient may now be positioned in supine position if tolerated

If aquatic option is available:

- Active assisted scapular plane elevation can be performed up to 90°
- Active assisted external rotation @ neutral can be performed up to 30°
- No resistance

Continue home exercise program:

- Grade 1-2 joint mobilizations to reduce pain, guarding, and tone
- PNF scapular movement patterns in side lying with elbow @ neutral
- Therapist guided passive mobility to 90° scapular elevation with grade 1-2 joint oscillations @ end range
- Continue isometrics
- Cervical and seated thoracic mobilizations as needed and guided by clinical examination / history.
- Continue week three exercises for elbow, wrist, and hand mobility
- Ice and modalities as needed
 - * No motoric response with electrical stimulation

WEEK FIVE:

Finalize incision healing progression

Assess patient compliance / understanding with home exercise program

Patient may be positioned in all positions (except directly on involved side) as tolerated

Only if aquatic option is available: Active scapular plane elevation to 90° with no resistance can be added to progressions

Continue home exercise program:

- Active assisted range of motion exercises in gravity eliminated positions as tolerated
- Scapular PNF movement patterns in varying positions
 - * Quadruped, side lying, standing, and seated
 - * No weight bearing on involved shoulder
- Therapist guided passive mobility to 120° scapular elevation as tolerated by patient's pain response.
- Continue isometrics
- Cervical and thoracic extension and rotation mobilizations as needed and guided by clinical exam / history
- Continue elbow, wrist, and hand mobility exercises
- Ice and modalities as needed
 - * No motoric response with electrical stimulation

SUMMARY:

Begin physical therapy at week three if instructed by surgeon.

Incision healing monitoring and compliance with home exercise program.

Focus on preventing inhibition and scapular control early in healing process.

Improve patient awareness of cervical and thoracic joint posture

Reduce postural musculature tone and desensitize incision site

PHASE TWO

Week 6-12 | Mobility

GOALS:

Discharge sling

Protect surgical repair by avoiding unwanted strain to the replacement

Begin to restore full active and passive mobility in the shoulder girdle

No strengthening of shoulder (above 2lb) until week 12

THINGS TO AVOID:

No pushing, pulling, or lifting anything larger than a cup of coffee (approx. 1-2 pounds). Avoid pushing yourself up from a chair, bed, or from your car.

While we encourage you to tease, touch, and nudge any pain you might experience, please don't push through the pain. Don't perform any activity that requires more range of motion than you comfortably have during this phase.

No movements that require excessive behind the back movements. Avoid jerky movements behind your back (putting wallet in pocket, putting belt on, or tucking in a shirt)

We will NOT begin running during this phase

THINGS TO BE DOING:

Try and go back to "normal life" as safely as possible. You can now use your shoulder for activities such as dressing, bathing, typing, grooming, and driving.

You will begin progressing your home exercises in physical therapy. We want patients to be doing their exercises 1-2x / day.

Physical therapy frequency will increase to 2-3x / week at this time.

Patient can continue to use ice machine for pain relief. However, heat may be added before therapy if needed.

WEEK 6 TO WEEK 12:

Joint mobilizations and manual therapy as needed to improve mobility and reduce pain and guarding

- Grade 1-4 mobilizations as needed
- Instrument assisted soft tissue mobilizations

Active and active assisted shoulder motion without restriction

- Forward elevation in scapular plane
- Side lying external rotation and abduction
- Supine shoulder flexion (active or active assisted)
- Ball rolls
- Wall / table slides
- Wand assisted shoulder movements
- Supine serratus elevations with circles
- Forward reaching (active or active assisted with cane)
- Recumbent bike

Passive shoulder motion without restriction

- Pulley apparatus (if proper joint arthrokinematics are noticed)
- Prayer stretch
- Doorway stretches (avoid 90/90 positions with anatomical shoulder replacements)
- Avoid internal rotation movements with reverse total shoulder replacement

Scapular stabilization and setting exercises

- PNF scapular movements in all positions
- Bilateral external rotation with manual cues for lower trap
- Scapular clocks
- Protraction and retraction with control and prolonged holds

Aquatic therapy for range of motion (if available). No resistance, just motion.

Regional Interdependence

- Cardiovascular demands. Begin increasing cardiovascular benefits with elliptical, stationary bike, or recumbent biking. Avoidance of running is instructed in this phase.
- Sagittal and frontal plane lunges
- Step ups
- Balance and lower extremity proprioceptive work

PHASE THREE

Week 12+ | Strength

GOALS:

Protect surgical repair until strength and mobility is full

Begin gradually restoring strength, power, and endurance in the shoulder girdle complex

Improve shoulder stability with initiating and progressing weight bearing activities in the shoulder

THINGS TO AVOID:

No sudden jerking or uncontrolled movements

No lifting objects away from the body that is heavier than 5 lbs.

No empty can (thumbs down) position with weights.

THINGS TO BE DOING:

Continue to use your shoulder as normal as possible for daily activities.

Strengthening exercises will be added to your home exercise regime. Patients should be performing their exercises 1x/day.

Physical therapy frequency can decrease to 1-2x / week if patient demonstrates good compliance with home exercise program.

Home exercise resistance equipment may include Thera bands, free weights, and body weight.

WEEK 12+

Gradual progression on rotator cuff strength with bands / free weights

- Thrower's Ten Program
 - * Advanced Thrower's Ten Program
- Body Blade
- Prone rowing with shoulder rotation progression
- Prone I's / Y's / and T's

Scapular strengthening progression and deltoid force coupling

- Three-way rowing
- Scapular push up with plus
- Bilateral external rotation with elevation
- Serratus wall slides

Closed-Kinetic chain upper extremity exercise progression

- Quadruped opposite arm / opposite leg
- Wall push ups
- Seated press ups

PNF shoulder movement patterns

- D1 and D2 flexion and extension in varying positions
 - * Standing, lunge, and supine

Regional Interdependence

- Bicep and triceps strengthening
- Progress lower extremity cardiovascular endurance
- Running may be progressed in this phase

Return to sport and work hardening activities begin at week 18

- Plyometric shoulder progression can be started at week 18

If the patient is an overhead or throwing athlete and you wish to contact us about our throwing or golf progressions after the physical therapy protocol, please feel free to contact us at:

UAMS Health
| Orthopaedics & Sports Medicine

UAMS Orthopaedics and Sports Medicine | 161 West Van Asche Drive, Fayetteville, AR 72703
479-966-4491 | uamshealth.edu